

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
20 March 2003 (20.03.2003)

PCT

(10) International Publication Number  
**WO 03/022236 A1**

- (51) International Patent Classification<sup>7</sup>: **A61K 7/48** 06611 (US). **ZHANG, Joanna, Hong**; Unilever Home & Personal Care USA, 40 Merritt Boulevard, Trumbull, CT 06611 (US).
- (21) International Application Number: PCT/EP02/09117
- (22) International Filing Date: 15 August 2002 (15.08.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/318,687 12 September 2001 (12.09.2001) US
- (71) Applicant (for AE, AG, AU, BB, BZ, CA, CY, GB, GD, GH, GM, IE, IL, KE, LC, LK, LS, MN, MW, NZ, OM, SD, SG, SL, SZ, TT, TZ, UG, VC, ZA, ZM, ZW only): **UNILEVER PLC** [GB/GB]; Unilever House, Blackfriars, London EC4P 4BQ (GB).
- (71) Applicant (for AL, AM, AT, AZ, BA, BE, BF, BG, BJ, BR, BY, CF, CG, CH, CI, CM, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, FR, GA, GE, GN, GQ, GR, GW, HR, HU, ID, IS, IT, JP, KG, KP, KR, KZ, LR, LT, LU, LV, MA, MC, MD, MG, MK, ML, MR, MX, MZ, NE, NL, NO, PH, PL, PT, RO, RU, SE, SI, SK, SN, TD, TG, TJ, TM, TN, TR, UA, UZ, VN, YU only): **UNILEVER NV** [NL/NL]; Weena 455, NL-3013 AL Rotterdam (NL).
- (71) Applicant (for IN only): **HINDUSTAN LEVER LIMITED** [IN/IN]; Hindustan Lever House, 165/166 Backbay Reclamation, Maharashtra, 400 020 Mumbai (IN).
- (72) Inventors: **SUARES, Alan, Joseph**; Unilever Home & Personal Care USA, 40 Merritt Boulevard, Trumbull, CT
- (81) Designated States (*national*): AE, AG, AL, AM, AT (utility model), AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ (utility model), CZ, DE (utility model), DE, DK (utility model), DK, DM, DZ, EC, EE (utility model), EE, ES, FI (utility model), FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK (utility model), SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**  
— with international search report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: THICKENED COSMETIC COMPOSITIONS COMPRISING A TAURATE COPOLYMER

(57) Abstract: A cosmetic composition is provided which includes an alpha- or beta-hydroxycarboxylic acid at least partially neutralized and a taurate copolymer in a low pH system. The taurate copolymer improves skin feel and provides viscosity to the system.



WO 03/022236 A1

- 1 -

## THICKENED COSMETIC COMPOSITIONS COMPRISING A TAURATE COPOLYMER

The invention relates to improved thickening systems for cosmetic compositions, particularly those in lotion and cream form.

Aqueous cosmetic compositions often require thickeners to achieve an aesthetically pleasing viscosity. Fluids that flow with a watery consistency too rapidly run off the treated skin areas. For a cosmetic to be effective, it often must have substantivity. Thickeners provide this substantivity. Furthermore, low viscosity formulas which may be skin effective nevertheless through their wateriness signal ineffectiveness to the consumer. Products of watery consistency are also aesthetically displeasing to consumers with expectations of rich and creamy products.

U.S. Patent 5,422,112 (Williams) discloses a triple thickener system including xanthan gum, magnesium aluminum silicate and polyacrylamide. The compositions are said to be particularly effective for thickening alpha-hydroxy carboxylic acids and salts thereof, especially at low pH.

U.S. Patent 5,874,095 (Deckner et al.) reports an enhanced skin penetration system for improved topical delivery of drugs. Essential to the system is a nonionic polyacrylamide of high molecular weight described as effective at low pH.

U.S. Patent 5,952,395 (Lorant) and U.S. Patent 5,891,452 (Sebillote-Arnaud et al.) describe cosmetic compositions gelled into an emulsion with a cross-linked poly(2-acrylamido-2-methylpropanesulfonic acid).

- 2 -

Countless numbers of other thickening agents are known in the literature. Perhaps this plethora intimates that not all thickening agents are equally effective for any particular type of formulation.

- 5 Indeed, there are some formulations which are extremely difficult to thicken, and even if initially thickened may have storage stability problems. Low pH systems are particularly sensitive and difficult.
- 10 Accordingly, it is an advantage of the present invention to provide a thickener system and thickened cosmetic compositions of sufficiently aesthetically pleasing viscosity and skinfeel.

It is another advantage of the present invention to provide  
15 thickening systems for cosmetic compositions that are effective at low pH.

It is still another advantage of the present invention to provide thickening systems for water and oil emulsion  
20 cosmetic compositions that also function as stabilizers preventing phase separation.

These and other advantages of the present invention will more readily become apparent from the description and examples which follow.

25 A cosmetic composition is provided which includes:

- (i) from 0.01 to 20% of a C<sub>1</sub>-C<sub>25</sub> alpha- or beta-hydroxycarboxylic acid at least partially present as a salt thereof;

- 3 -


- (ii) from 0.01 to 10% of a taurate copolymer; and
- (iii) a cosmetically acceptable carrier, wherein the composition has a pH of less than 6.

5 Now it has been discovered that taurate copolymers are highly effective thickening agents for low pH cosmetic compositions. This system is particularly useful for building viscosity in relatively acidic compositions, especially those containing C<sub>1</sub>-C<sub>25</sub> alpha- or beta-  
10 hydroxycarboxylic acids. Beyond building viscosity, the thickening system of this invention have the further advantage of stabilizing oil and water emulsions and providing a good skin feel.

Accordingly, a first element of compositions according to  
15 the present invention is that of a taurate copolymer. A particularly preferred copolymer is one wherein the taurate repeating monomer unit is acryloyl dimethyl taurate (in either free acid or salt form). Monomers forming the copolymer with taurate may include: styrene, acrylic acid,  
20 methacrylic acid, vinyl chloride, vinyl acetate, vinyl pyrrolidone, isoprene, vinyl alcohol, vinyl methylether, chloro-styrene, dialkylamino-styrene, maleic acid, acrylamide, methacrylamide and mixtures thereof. Where the term "acid" appears, the term means not only the free acid  
25 but also C<sub>1</sub>-C<sub>30</sub> alkyl esters, anhydrides and salts thereof. Preferably but not exclusively the salts may be ammonium, alkanolammonium, alkali metal and alkaline earth metal salts. Most preferred are the ammonium and alkanolammonium salts.

- 4 -

Most preferred as the copolymer is Acryloyl Dimethyltaurate/Vinyl Pyrrolidone Copolymer, which is the INCI nomenclature, for a material supplied by Clariant Corporation under the trademark Aristoflex<sup>®</sup> AVC, having the following general formula:




wherein n and m are integers which may independently vary from 1 to 10,000.

Average molecular weight of copolymers according to the invention may range from 1,000 to 3,000,000, preferably from 3,000 to 100,000, optimally from 10,000 to 80,000.

Amounts of the taurate copolymer may range from 0.001 to 10%, preferably from 0.01 to 8%, more preferably from 0.1 to 5%, optimally from 0.2 to 1% by weight of the composition.

A second element of the compositions according to the present invention is an alpha- or beta-hydroxycarboxylic acid or salt thereof. The former may be a C<sub>1</sub>-C<sub>25</sub> alpha-hydroxycarboxylic acid of formula I

- 5 -



wherein R and R<sup>1</sup> are independently H, F, Cl, Br, alkyl,  
 aralkyl or aryl groups being saturated or unsaturated,  
 5 isomeric or nonisomeric, straight or branched chain, or in  
 cyclic form having 5 or 6 ring members, and in addition, R  
 and R<sup>1</sup> may carry OH, CHO, COOH and alkoxy groups having 1 to  
 9 carbon atoms, the  $\alpha$ -hydroxyacid existing as a free acid or  
 lactone form, or in salt form with an organic amine base or  
 10 an inorganic alkali, and as stereoisomers, and D, L, and DL  
 forms when R and R<sup>1</sup> are not identical.

Illustrative of this group of materials are 2-  
 hydroxyethanoic acid (glycolic acid); 2-hydroxypropanoic  
 15 acid (lactic acid); 2-methyl 2-hydroxypropanoic acid  
 (methyllactic acid); 2-hydroxybutanoic acid; 2-  
 hydroxypentanoic acid; 2-hydroxyhexanoic acid; 2-  
 hydroxyheptanoic acid; 2-hydroxyoctanoic acid; 2-  
 hydroxynonanoic acid; 2-hydroxydecanoic acid; 2-  
 20 hydroxyundecanoic acid; 2-hydroxydodecanoic acid ( $\alpha$ -  
 hydroxylauric acid); 2-hydroxytetradecanoic acid ( $\alpha$ -  
 hydroxymyristic acid); 2-hydroxyhexadecanoic acid ( $\alpha$ -  
 hydroxypalmitic acid); 2-hydroxyoctadecanoic acid ( $\alpha$ -  
 hydroxystearic acid); 2-hydroxyeicosanoic acid ( $\alpha$ -  
 25 hydroxyarachidonic acid); 2-phenyl 2-hydroxyethanoic acid

- 6 -

(mandelic acid); 2,2-diphenyl 2-hydroxyethanoic acid  
(benzilic acid); 3-phenyl 2-hydroxypropanoic acid  
(phenyllactic acid); 2-phenyl 2-methyl 2-hydroxyethanoic  
acid (atrolactic acid); 2-(4'-hydroxyphenyl) 2-  
5 hydroxyethanoic acid; 2-(4'-chlorophenyl 2-hydroxyethanoic  
acid; 2-(3'-hydroxy-4'-methoxyphenyl) 2-hydroxyethanoic  
acid; 2-(4'-hydroxy-3'-methoxyphenyl) 2-hydroxyethanoic  
acid; 3-(2-hydroxyphenyl) 2-hydroxypropanoic acid; 3-(4'-  
hydroxyphenyl) 2-hydroxypropanoic acid; and 2-(3',4'-  
10 dihydroxyphenyl) 2-hydroxyethanoic acid.

Most preferred of this group of materials are glycolic acid,  
lactic acid, and 2-hydroxyoctanoic acids. The salts may be  
selected from alkalimetal, ammonium and C<sub>1</sub>-C<sub>20</sub> alkyl or  
15 alkanolammonium counterions.

Advantageously, the hydroxycarboxylic acid will be fully  
neutralized as the potassium, sodium, ammonium or  
alkanolammonium salt. At least 1%, preferably at least 10%,  
and optimally at least 20% will be present as the salt form.  
20 Also effective are mixtures of acid and salt forms ranging  
in molar ratio from 1000:1 to 1:1000, preferably from 100:1  
to 1:100, more preferably from 50:1 to 1:1, optimally from  
20:1 to 2:1.

The beta-hydroxycarboxylic acids are best illustrated by  
25 salicylic acid and its derivatives. Levels of the  
hydroxycarboxylic acids may range from 0.01 to 20%,  
preferably from 0.2 to 10%, optimally from 1 to 5% by  
weight.

- 7 -

Compositions of the present invention may either be aqueous or anhydrous. Preferably the compositions are aqueous, especially water and oil emulsions of the W/O or O/W variety. Water when present will be in amounts which may range from 5  
5 to 90%, preferably from 35 to 70%, optimally between 40 and 60% by weight.

The pH of compositions of this invention is less than 7. Advantageously, pH may range from 1 to 6, preferably from 2 to 5, optimally from 2.5 to 3.8.

10 Emollient materials in the form of mineral oils, silicone oils and synthetic esters may be incorporated into compositions of the present invention. Amounts of the emollients if incorporated may range anywhere from 0.1 to 30%, preferably between 0.5 and 20% by weight.

15 Silicone oils may be divided into the volatile and non-volatile variety. The term "volatile" as used herein refers to those materials which have a measurable vapor pressure at ambient temperature. Volatile silicone oils are preferably chosen from cyclic or linear polydimethylsiloxanes  
20 containing from 3 to 9, preferably from 4 to 5, silicon atoms.

Linear volatile silicone materials generally have viscosities less than about 5 centistokes at 25°C while cyclic materials typically have viscosities of less than  
25 about 10 centistokes.

Nonvolatile silicone oils useful as an emollient material include polyalkyl siloxanes, polyalkylaryl siloxanes and polyether siloxane copolymers. The essentially non-volatile



- 8 -

polyalkyl siloxanes useful herein include, for example, polydimethyl siloxanes with viscosities of from about 5 to about 100,000 centistokes at 25°C.

Among suitable ester emollients are:

5

(1) Alkenyl or alkyl esters of fatty acids having 10 to 20 carbon atoms. Examples thereof include isopropyl palmitate, isopropyl isostearate, isononyl isonanonate, oleyl myristate, oleyl stearate, and oleyl oleate.

10

(2) Ether-esters such as fatty acid esters of ethoxylated fatty alcohols.

15

(3) Polyhydric alcohol esters. Ethylene glycol mono and di-fatty acid esters, diethylene glycol mono- and di-fatty acid esters, polyethylene glycol (200-6000) mono- and di-fatty acid esters, propylene glycol mono- and di-fatty acid esters, polypropylene glycol 2000 monooleate, polypropylene glycol 2000 monostearate, ethoxylated propylene glycol monostearate, glyceryl mono- and di-fatty acid esters, polyglycerol poly-fatty esters, ethoxylated glyceryl monostearate, 1,3-butylene glycol monostearate, 1,3-butylene glycol distearate, polyoxyethylene polyol fatty acid ester, sorbitan fatty acid esters, and polyoxyethylene sorbitan fatty acid esters are satisfactory polyhydric alcohol esters.

20

25

30

- 9 -

(4) Wax esters such as beeswax, spermaceti, myristyl myristate, stearyl stearate.

(5) Sterols esters, of which soya sterol and cholesterol fatty acid esters are examples thereof.

The most preferred esters are dicaprylyl ether and isopropyl isostearate.

Fatty acids having from 10 to 30 carbon atoms may also be included in the compositions of this invention. Illustrative of this category are pelargonic, lauric, myristic, palmitic, stearic, isostearic, hydroxystearic, oleic, linoleic, ricinoleic, arachidic, behenic and erucic acids.

Humectants of the polyhydric alcohol-type may also be included in the compositions of this invention. The humectant aids in increasing the effectiveness of the emollient, reduces scaling, stimulates removal of built-up scale and improves skin feel. Typical polyhydric alcohols include glycerol (also known as glycerin), polyalkylene glycols and more preferably alkylene polyols and their derivatives, including propylene glycol, dipropylene glycol, polypropylene glycol, polyethylene glycol and derivatives thereof, sorbitol, hydroxypropyl sorbitol, hexylene glycol, 1,3-butylene glycol, 1,2,6-hexanetriol, ethoxylated glycerol, propoxylated glycerol and mixtures thereof. For best results the humectant is preferably glycerin. The amount of humectant if present may range anywhere from 0.5 to 30%, preferably between 1 and 15% by weight of the composition.

Collectively the water, solvents, silicones, esters, fatty acids and/or humectants are viewed as cosmetically acceptable

- 10 -

carriers for the compositions of the invention. Total amount of carrier will typically range from 1 to 99.9%, preferably from 80 to 99% by weight.

Cosmetic compositions of the present invention may be in any  
5 form. These forms may include lotions, creams, roll-on formulations, mousses, aerosol sprays and cloth- or pad-applied formulations.

Emulsifiers may also be present in cosmetic compositions of the present invention. Total concentration of the emulsifier  
10 if present will range from 0.1 to 40%, preferably from 1 to 20%, optimally from 1 to 5% by weight of the total composition. The emulsifier may be selected from the group consisting of anionic, nonionic, cationic and amphoteric actives. Particularly preferred nonionic surfactants are  
15 those with a C<sub>10</sub>-C<sub>20</sub> fatty alcohol or acid hydrophobe condensed with from 2 to 100 moles of ethylene oxide or propylene oxide per mole of hydrophobe; C<sub>2</sub>-C<sub>10</sub> alkyl phenols condensed with from 2 to 20 moles of alkylene oxide; mono- and di- fatty acid esters of ethylene glycol; fatty acid  
20 monoglyceride; sorbitan, mono- and di- C<sub>8</sub>-C<sub>20</sub> fatty acids; and polyoxyethylene sorbitan as well as combinations thereof. Alkyl polyglycosides and saccharide fatty amides (e.g. methyl gluconamides) are also suitable nonionic emulsifiers.

Preferred anionic emulsifiers include soap, alkyl ether  
25 sulfate and sulfonates, alkyl sulfates and sulfonates, alkylbenzene sulfonates, alkyl and dialkyl sulfosuccinates, C<sub>8</sub>-C<sub>20</sub> acyl isethionates, C<sub>8</sub>-C<sub>20</sub> alkyl ether phosphates, alkylethercarboxylates and combinations thereof.

- 11 -

Sunscreen actives may also be included in compositions of the present invention. Particularly preferred are such materials as Avobenzene, available as Parsol<sup>®</sup> 1789, ethylhexyl p-methoxycinnamate, available as Parsol<sup>®</sup> MCX, and benzophenone-  
5 3, also known as Oxybenzone. Inorganic sunscreen actives may be employed such as microfine titanium dioxide, polyethylene and various other polymers. Amounts of the sunscreen agents if present will generally range from 0.1 to 30%, preferably from 2 to 20%, optimally from 4 to 10% by weight.

10 Preservatives can desirably be incorporated into the cosmetic compositions of this invention to protect against the growth of potentially harmful microorganisms. Suitable traditional preservatives for compositions of this invention are alkyl esters of para-hydroxybenzoic acid. Other preservatives which  
15 have more recently come into use include hydantoin derivatives, propionate salts, and a variety of quaternary ammonium compounds. Cosmetic chemists are familiar with appropriate preservatives and routinely choose them to satisfy the preservative challenge test and to provide product  
20 stability. Particularly preferred preservatives are iodopropynyl butyl carbamate, phenoxyethanol, methyl paraben, propyl paraben, imidazolidinyl urea, sodium dehydroacetate and benzyl alcohol. The preservatives should be selected having regard for the use of the composition and possible  
25 incompatibilities between the preservatives and other ingredients in the emulsion. Preservatives are preferably employed in amounts ranging from 0.01% to 2% by weight of the composition.

- 12 -

Co-thickening agents may be included in compositions of the present invention. Particularly useful are the polysaccharides. Examples include starches, natural/synthetic gums and cellulosics. Representative of the starches are  
5 chemically modified starches such as aluminum starch octenylsuccinate. Suitable gums include xanthan, sclerotium, pectin, karaya, arabic, agar, guar, carrageenan, alginate and combinations thereof. Suitable cellulosics include  
10 hydroxypropyl cellulose, hydroxypropyl methylcellulose, ethylcellulose and sodium carboxy methylcellulose. Most preferred is aluminum starch octenylsuccinate.

Amounts of the co-thickener may range from 0.001 to 5%, preferably from 0.1 to 2%, optimally from 0.2 to 0.5% by weight.

15 For additional thickening, it is preferred to have magnesium aluminum silicate, commercially available as Veegum<sup>®</sup>, sold by the R.T. Vanderbilt Company. Amounts of this inorganic thickening agent if present may range from 0.01 to 10%, preferably from 0.5 to 1.2% by weight.

20 Minor adjunct ingredients may also be present in the cosmetic compositions. Among them may be the water-insoluble vitamins such as Vitamin A Palmitate, Vitamin E Acetate and DL-panthenol. Also useful are: retinol, ceramides and herbal extracts including green tea and chamomile.

25 Colorants, fragrances and abrasives may also be included in compositions of the present invention. Each of these substances may range from 0.05 to 5%, preferably between 0.1 and 3% by weight.

- 13 -

Compositions of the present invention advantageously will have viscosities ranging from 500 to 100,000, preferably from 1,000 to 60,000, optimally from 1,500 to 30,000 cps. These viscosities are measured on a Brookfield RVT

5 Viscometer, Spindle 4, at 20 rpm and 23°C.

Except in the operating and comparative examples, or where otherwise explicitly indicated, all numbers in this description indicating amounts of material ought to be understood as modified by the word "about".

10 The term "comprising" is meant not to be limiting to any subsequently stated elements but rather to encompass non-specified elements of major or minor functional importance. In other words the listed steps, elements or options need not be exhaustive. Whenever the words "including" or  
15 "having" are used, these terms are meant to be equivalent to "comprising" as defined above.

The following Examples will more fully illustrate the embodiments of this invention. All parts, percentages and proportions referred to herein and in the appended claims are  
20 by weight unless otherwise indicated.

#### EXAMPLES 1-8

Typical formulations according to the present invention are described below.

[illegible]

- 15 -

## EXAMPLE 9

A series of comparative experiments were conducted to measure the relative thickening effectiveness of various polymers in low pH systems containing hydroxycarboxylic acid. Table 1 reports a series of viscosity measurements on a gel solution consisting of 98% water and 2% thickener. Measurements were conducted using a Brookfield RVT instrument, Spindle No. 4 at 20 rpm and 23°C. The most efficient thickener was Aristoflex<sup>®</sup> AVC (Sample 1G).

10

TABLE 1\*

Sample	Thickener	Viscosity (cps)	pH
1A	Sepigel <sup>®</sup> 305	19,995	6.0
1B	Simulgel <sup>®</sup> EG	23,900	6.8
1C	Simulgel <sup>®</sup> NS	16,600	5.5
1D	Stabylen <sup>®</sup> 30	15,850	2.9
1E	Synthalen <sup>®</sup> CR	27,750	4.0
1F	Carbopol <sup>®</sup> 934	1,735	2.8
1G	Aristoflex <sup>®</sup> AVC	40,000	4.8

\* 2% Gel



- 16 -

The identities of the polymers in Table 1 are as follows:

Thickener	INCI Name
Sepigel® 305	Polyacrylamide/C13-14 Isoparaffin/Laureth-7
Simulgel® EG	Sodium Acrylate/Sodium Acryloyldimethyl Taurate Copolymer/Isohexadecane/Polysorbate 80
Simulgel® NS	Hydroxyethyl Acrylate/Sodium Acryloyldimethyl Taurate Copolymer/Squalane/Polysorbate 60
Stabylen® 30	Acrylates/Vinyl Isodecanoate Crosspolymer
Synthalen® CR	Polyquaternium-37 (Quaternary Acrylic Polymer)
Carbopol® 934	Crosslinked Polyacrylate
Aristoflex® AVC	Acryloyl Taurate/Vinyl Pyrrolidone Copolymer

Table 2 reports the effect of adding glycolic acid (an alpha-hydroxycarboxylic acid) to each of the gel solutions. All of the thickened samples decreased in viscosity. However, the Synthalen® CR and Aristoflex® AVC samples still maintained a reasonable level of thickness.

TABLE 2\*

Sample	Thickener	Viscosity (cps)	pH
1A	Sepigel® 305	<1,000	2.0
1B	Simulgel® EG	<100	2.1
1C	Simulgel® NS	<100	2.0
1D	Stabylen® 30	<100	1.9
1E	Synthalen® CR	17,450	1.7
1F	Carbopol® 934	<100	1.9
1G	Aristoflex® AVC	20,500	2.0

\* 2% Gel + 8% Glycolic Acid

Tables 3 and 4 illustrates the effect of partially neutralizing the glycolic acid with ammonium hydroxide to result in a partially neutralized acid solution of glycolic

- 17 -

acid/ammonium glycolate. In these highly stressed, low pH systems, only the Aristoflex<sup>®</sup> AVC provided even a modest measure of thickening to the hydroxycarboxylic acid systems.

TABLE 3\*

Sample	Thickener	Viscosity (cps)	PH
1A	Sepigel <sup>®</sup> 305	<1,00	3.0
1B	Simulgel <sup>®</sup> EG	<100	3.0
1C	Simulgel <sup>®</sup> NS	<100	3.0
1D	Stabylen <sup>®</sup> 30	<100	3.0
1E	Synthalen <sup>®</sup> CR	3,650	2.9
1F	Carbopol <sup>®</sup> 934	<100	3.0
1G	Aristoflex <sup>®</sup> AVC	9,060	3.0

5 \* 2% Gel + 8% Glycolic Acid/Ammonium Glycolate (acid to salt molar ratio of 10:1)

TABLE 4\*

Sample	Thickener	Viscosity (cps)	PH
1A	Sepigel <sup>®</sup> 305	<100	3.6
1B	Simulgel <sup>®</sup> EG	<100	3.6
1C	Simulgel <sup>®</sup> NS	<100	3.5
1D	Stabylen <sup>®</sup> 30	1,445	3.5
1E	Synthalen <sup>®</sup> CR	<100	3.5
1F	Carbopol <sup>®</sup> 934	<100	3.5
1G	Aristoflex <sup>®</sup> AVC	2,555	3.6

\* 2% Gel + 8% Glycolic Acid/Ammonium Glycolate (acid to salt molar ratio of 3.4:1)

- 18 -

CLAIMS

1. A cosmetic composition comprising:
  - (i) from 0.01 to 20% of a C<sub>1</sub>-C<sub>25</sub> alpha- or beta-hydroxy  
5 carboxylic acid at least partially present as a  
salt thereof;
  - (ii) from 0.01 to 10% of a taurate copolymer; and
  - (iii) a cosmetically acceptable carrier, wherein the  
composition has a pH of less than 7.
- 10 2. A composition according to claim 1 wherein the C<sub>2</sub>-C<sub>25</sub>  
alpha-hydroxy carboxylic acid is selected from glycolic  
acid, lactic acid, 2-hydroxyoctanoic acid and  
combinations thereof.
- 15 3. A composition according to claim 1 or claim 2 wherein  
the taurate copolymer is at least partially formed from  
acryloyl dimethyltaurate as a monomer unit.
- 20 4. A composition according to any of the preceding claims  
wherein the hydroxycarboxylic acid and salt thereof are  
present in a molar ratio of 1000:1 to 1:1000.
- 25 5. A composition according to claim 4 wherein the  
hydroxycarboxylic acid and salt thereof are present in a  
molar ratio of 50:1 to 1:1.
- 30 6. A composition according to any of claims 3 to 5 wherein  
the acryloyl dimethyltaurate is copolymerized with a  
monomer selected from the group consisting of styrene,

- 19 -

acrylic acid, methacrylic acid, vinyl chloride, vinyl acetate, vinyl pyrrolidone, isoprene, vinyl alcohol, vinyl methylether, chloro-styrene, maleic acid, acrylamide, methacrylamide and mixtures thereof.

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 02/09117

**A. CLASSIFICATION OF SUBJECT MATTER**  
IPC 7... A61K7/48

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, PAJ, WPI Data

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 815 845 A (OREAL) 7 January 1998 (1998-01-07) claims 1-3,17,18 examples 1,4,6 page 2, line 17 - line 54 ----	1-6
X	WO 01 28338 A (PROCTER & GAMBLE) 26 April 2001 (2001-04-26) claim 1 example 3 page 41, line 5 - line 9 ----	1-6
X	WO 01 62214 A (COLOR ACCESS INC) 30 August 2001 (2001-08-30) claims 1-3 example 1 page 2, line 11 -page 3, line 14 ----- -/--	1-6

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*&\* document member of the same patent family

Date of the actual completion of the international search

21 November 2002

Date of mailing of the international search report

28/11/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Simon, F

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 02/09117

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>US 5 863 545 A (GRIAT JACQUELINE)  26 January 1999 (1999-01-26)  claim 1  examples 1-3  column 2, line 3 - line 44  -----</p>	1-6

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 02/09117

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0815845	A	07-01-1998	FR 2750326 A1	02-01-1998
			AT 189117 T	15-02-2000
			BR 9702539 A	29-09-1998
			CA 2209430 A1	28-12-1997
			DE 69701205 D1	02-03-2000
			DE 69701205 T2	31-05-2000
			EP 0815845 A1	07-01-1998
			ES 2144831 T3	16-06-2000
			JP 2941234 B2	25-08-1999
			JP 10067616 A	10-03-1998
			KR 251980 B1	01-07-2000
			PL 320776 A1	05-01-1998
			RU 2167642 C2	27-05-2001
			US 6468549 B1	22-10-2002
WO 0128338	A	26-04-2001	US 6294186 B1	25-09-2001
			AU 1095901 A	30-04-2001
			AU 8029400 A	30-04-2001
			AU 8029800 A	30-04-2001
			BR 0014776 A	16-07-2002
			BR 0014905 A	18-06-2002
			BR 0014906 A	11-06-2002
			EP 1221852 A2	17-07-2002
			EP 1221851 A2	17-07-2002
			EP 1238159 A1	11-09-2002
			WO 0128338 A2	26-04-2001
			WO 0129315 A1	26-04-2001
			WO 0128337 A2	26-04-2001
			US 6475501 B1	05-11-2002
WO 0162214	A	30-08-2001	AU 3707001 A	03-09-2001
			CN 1362873 T	07-08-2002
			EP 1189579 A1	27-03-2002
			WO 0162214 A1	30-08-2001
			US 2002058055 A1	16-05-2002
US 5863545	A	26-01-1999	FR 2709982 A1	24-03-1995
			US 6001379 A	14-12-1999
			AT 175864 T	15-02-1999
			CA 2132144 A1	16-03-1995
			DE 69416060 D1	04-03-1999
			DE 69416060 T2	27-05-1999
			EP 0642781 A1	15-03-1995
			ES 2129602 T3	16-06-1999
			JP 2693116 B2	24-12-1997
			JP 7163859 A	27-06-1995
			US 5531993 A	02-07-1996